


8. SYMMETRY-BREAKING TECHNIQUES

Closed-shell vs open-shell nuclei:


Mean-field configurations have different features as N and Z vary. In particular, one distinguishes between closed- and open-shell nuclei:

CLOSED SHELL



nucleons entirely fill levels below a magic number

OPEN SHELL



nucleons partially fill levels below a magic number

- large energy gap at the Fermi surface \Rightarrow excitations hindered \Rightarrow very stable configurations
- dynamical correlations dominate

- small energy gap (or $\rightarrow 0$) at the Fermi surface \Rightarrow excitations favoured \Rightarrow less stable configurations
- static properties essential

$$\Delta E_{\text{HBPT}}^{(2)} \propto \frac{1}{e_a + e_b - e_i - e_j}$$


Well-defined particle-hole expansion

Expansion breaks down (diverges or even ill-defined)

Possible solution: use symmetry breaking (and symmetry restoration afterwards) to capture strong static correlations.

Symmetries

In nuclear physics $[\hat{X}, H] = 0$ for



The solutions $|\Psi_i^x\rangle$ of the many-body Schrödinger equation

$$H |\Psi_i^x\rangle = E_i |\Psi_i^x\rangle$$

are labeled by quantum numbers $\{x\}$:

- $N, Z \in \mathbb{N}$
- $\vec{P} \in \mathbb{R}^3$
- $2J \in \mathbb{N}$ and $2M \in \mathbb{Z}$ such that $-J \leq M \leq J$
- $\Pi = \pm 1$
- $T^2 = (-1)^N (-1)^Z$

The nuclear part of H , in addition, nearly commutes with the isospin operators \hat{T}^2, \hat{T}_Z .

In which space do we vary $|\phi\rangle$ (or $\{\psi_a\}$ or ρ_{op}) (2)
when minimising $E[\rho]$?

The natural way is to perform the variation such that $|\phi\rangle$ carries the same quantum numbers $\{x\}$ of $|\psi_0^x\rangle$.

However, this might be too constrained for a relatively simple wave function like a Slater determinant!

Example:

$$\hat{P}|\phi\rangle = \vec{P}|\phi\rangle$$

$$\Rightarrow \psi(\vec{r}') = \psi_{\vec{k}}(\vec{r}') = e^{i\vec{k}\cdot\vec{r}'}$$

$$\Rightarrow \rho(\vec{r}') = \rho \quad \text{constant density}$$

↓

Misses the correlations that induce spatial localisation of the interelectron motion

→ usually broken in nuclear structure calculations

→ similar situation holds for other symmetries

Spontaneous symmetry breaking

Expand the horizon: the variation of $|\phi\rangle$ can be performed allowing for (spontaneous) symmetry breaking.

Let $|\phi\rangle$ span different irreducible representations (IRREPs) of a symmetry group, e.g. $SU(2)$

$$|\phi\rangle = \sum_{J=0,2,4,\dots} \sum_{M \leq |J|} c_{JM} |O^{JM}\rangle$$

(3)

For an even-even nucleus, the ground state is characterised by $J=0$. Here one chooses to consider a linear combination of $J=0$ and higher (even) values of J .

By doing this:

- Certain (long-range) correlations are included in $|\phi\rangle$. As a consequence, the corresponding energy \bar{E}^{HF} will be generally lower than for a symmetry-conserving variation.
- However, $|\phi\rangle$ may not carry good quantum numbers. One can see this as describing a wave packet rather than the ground state. It might be problematic to compute certain quantities, e.g. those relying on selection rules (e.g. $B\bar{E}(2)$).
- Symmetries have to be eventually restored: projected HF. This typically brings additional correlation energy.
- Note that the shell model follows an approach orthogonal to this and expand

$$|\Psi_0^x\rangle = \sum_k c_{0k} |\Phi_k^x\rangle$$

— symmetry-conserving excited Slater determinants

(1)

Physical content of symmetry breaking

Symmetry	Group	Casimir	Correlations	Type of nuclei	Excitation pattern
Translation	$T(3)$	\hat{p}^2	Spatial localisation	All	Surface vibrations
Rotations in iso space	$SU(2)$	\hat{J}^2	angular localisation	Doubly open shell	Rotational bands
Rotations in gauge space	$U(1)$	\hat{N}, \hat{E}	pairing (superfluidity)	Singly and doubly open shell	Energy gap

• Symmetries can be enforced or relaxed (in the variation of $|\psi\rangle$) depending on the nucleus

• Enforcing symmetries allows to exploit them in the calculations
 → work with an effective basis
 → reduced dimensionality of the problem and gain in CPU costs

HARTREE-FOCK - BOGOLYUBOV APPROXIMATION

(5)

Bogolyubov states

Consider creation operators $a_\alpha^+ |0\rangle = |\alpha\rangle$ with $\{a_\alpha, a_\beta^+\} = \delta_{\alpha\beta}$

Slater determinant

- $|\phi\rangle \equiv \prod_{i=1}^A b_i^+ |0\rangle$

with

$$b_\mu^+ = \sum_\alpha U_{\alpha\mu} a_\alpha^+$$

unitary transformation



- vacuum state

$$\begin{cases} b_\mu^+ |\phi\rangle = 0 & \text{for } \mu \in [1, A] \\ b_\mu |\phi\rangle = 0 & \text{for } \mu \in [A, \infty] \end{cases}$$

- symmetry-conserving

$$\hat{A} |\phi\rangle = A |\phi\rangle$$

Bogolyubov state

- $|\phi\rangle \equiv \prod_\mu \beta_\mu |0\rangle$

with

$$\beta_\mu \equiv \sum_\alpha \left[U_{\alpha\mu}^* a_\alpha + V_{\alpha\mu}^* a_\alpha^+ \right]$$

quasiparticle operators

unitary transformation

$$\begin{pmatrix} U^\dagger & V^\dagger \\ V^\dagger & U^\dagger \end{pmatrix}$$



- vacuum state

$$\beta_\mu |\phi\rangle = 0 \quad \forall \mu$$

- symmetry-breaking

$$\hat{A} |\phi\rangle \neq A |\phi\rangle$$

Density matrices

(6)

In addition to the (hermitian) normal density matrix

$$\rho_{\alpha\beta} \equiv \langle \phi | b_{\beta}^{\dagger} b_{\alpha} | \phi \rangle = \rho_{\beta\alpha}^*$$

an (antisymmetric) anomalous density matrix appears

$$\kappa_{\alpha\beta} \equiv \langle \phi | b_{\beta} b_{\alpha} | \phi \rangle = -\kappa_{\beta\alpha}$$

encodes the symmetry breaking

If $\kappa \neq 0 \Rightarrow \rho^2 \neq \rho$ and one needs to introduce a generalised density matrix

$$R = \begin{pmatrix} \rho & \kappa \\ -\kappa^* & 1 - \rho^* \end{pmatrix}$$

that is now idempotent $R^2 = R$.

HFB energy

The variational space is enlarged to the full Fock space \mathcal{F}_N while keeping a simple product trial-state

$$|\phi\rangle \equiv \prod_{\mu} \beta_{\mu} |0\rangle$$

where β_{μ} mix b_{α} and b_{α}^{\dagger} but still fulfill standard anticommutation relations.

Wick's theorem with respect to $|\phi\rangle$ applied to H

(7)

gives

$$\begin{aligned} E^{HF3} &= \sum_{\alpha\beta} t_{\alpha\beta} p_{\beta\alpha} + \frac{1}{2} \sum_{\alpha\beta\gamma\delta} \bar{v}_{\alpha\beta\gamma\delta} p_{\gamma\alpha} p_{\delta\beta} + \frac{1}{4} \sum_{\alpha\beta\gamma\delta} \bar{v}_{\alpha\beta\gamma\delta} \kappa_{\alpha\beta}^* \kappa_{\gamma\delta} \\ &= E[p, \kappa, \kappa^*] \end{aligned}$$

Equations of motion

In the minimisation procedure, independent variables are

$$p_{ij}, p_{ij}^*, \kappa_{ij}, \kappa_{ij}^* \quad \text{for } j \leq i$$

The constraint $p^2 = p$ is generalised to $R^2 = R$

One needs to further constrain $\langle \phi | \hat{A} | \phi \rangle = A$ since a free variation would lead to

$$A \rightarrow \infty$$

$$E^{HF3} \rightarrow -\infty$$

↓
Hence set $\delta \left[E[p, \kappa, \kappa^*] - \lambda \text{Tr}\{p\} - \text{Tr}\{\Lambda (R^2 - R)\} \right] = 0$

"chemical potential"

matrix
of Lagrange parameters

under variations δR and get

(8)

$$\mathcal{H} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} h - \lambda & \Delta \\ -\Delta^* & -h^* + \lambda \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = E \begin{pmatrix} u \\ v \end{pmatrix}$$

where

$$h_{\alpha\beta} \equiv \frac{\delta E[\rho, \kappa, \kappa^*]}{\delta \rho_{\alpha\beta}} = t_{\alpha\beta} + \sum_{\beta\delta} \bar{V}_{\alpha\beta\delta\delta} \rho_{\delta\beta}$$

Hartree-Fock field

$$\Delta_{\alpha\beta} \equiv \frac{\delta E[\rho, \kappa, \kappa^*]}{\delta \kappa_{\alpha\beta}^*} = \frac{1}{2} \sum_{\beta\delta} \bar{V}_{\alpha\beta\delta\delta} \kappa_{\beta\delta}$$

Bogolyubov field (or pairing field)

- As for HF, this is not a standard eigenvalue problem since

$$\mathcal{H} = \mathcal{H}[\rho, \kappa, \kappa^*] = \mathcal{H}[u, v]$$

Correlations beyond HFB

(9)

Using Wick's theorem with respect to $|\phi\rangle$ in HFB quasi-particle basis

$$H = \underbrace{\bar{E}^{\text{HFB}} + \sum_{\alpha} E_{\alpha} \beta_{\alpha}^{\dagger} \beta_{\alpha}}_{H_0'} + \underbrace{:V_2:}_{H_1'}$$

HFB approximation consist in taking $H^{\text{HFB}} = H_0'$ and neglecting H_1'

Then $H^{\text{HFB}} |\phi\rangle = \bar{E}^{\text{HFB}} |\phi\rangle$

However H^{HFB} displays other eigenstates depicted as two-quasi-particle excitations on top of $|\phi\rangle$

$$H^{\text{HFB}} |\phi^{\alpha\beta\dots}\rangle = \bar{E}^{\alpha\beta\dots} |\phi^{\alpha\beta\dots}\rangle$$

with

$$\begin{cases} |\phi^{\alpha\beta\dots}\rangle = \beta_{\alpha}^{\dagger} \beta_{\beta}^{\dagger} \dots |\phi\rangle \\ \bar{E}^{\alpha\beta\dots} = \bar{E}^{\text{HFB}} + \bar{E}_{\alpha} + \bar{E}_{\beta} + \dots \end{cases}$$

The exact ground-state energy will be

$$\bar{E}_0 = \bar{E}^{\text{HFB}} + \Delta \bar{E}_0^{\text{HFB}}$$

correlation energy due to effects of H_1'

Brillouin's theorem states that $\langle \phi | H_1' | \phi^{\alpha\beta} \rangle = 0$

however H_1 couples $|\phi\rangle$ to $|\phi^{\alpha\beta\gamma\delta}\rangle, |\phi^{\alpha\beta\gamma\delta, \epsilon}\rangle, \dots$

(10)

i.e.

$$\langle \phi | H_1 | \phi^{\alpha\beta\gamma\delta} \rangle \neq 0$$

↓

Correlation expansion methods

Express exact wave function as

$$|\Psi\rangle = W |\phi\rangle \quad \text{reference state (HF or HFBS)}$$

Wave operator

(includes corrections on top of a reference state)

E.g. for HFBS reference state

$$|\Psi\rangle = |\phi\rangle + |\phi^{\alpha\beta}\rangle + |\phi^{\alpha\beta\gamma\delta}\rangle + \dots$$

Total energy is

$$E_0 = \langle \Psi | H | \Psi \rangle = \underbrace{\langle \phi | H_0 | \phi \rangle}_{E_{\text{HFBS}}} + \underbrace{\langle \phi | H_1 | \phi^{\alpha\beta} \rangle + \dots}_{\Delta E_0^{\text{HFBS}}}$$

scope of many-body techniques

Alternatively to ab initio calculations, there exist more "effective" approaches.

Two main possibilities to reduce the complexity of

$$H |\Psi\rangle = E |\Psi\rangle$$

Reduce 1-body Hilbert space



Work in a subspace ("valence space") with a reduced number of nucleons



(Interacting) shell model approach

Reduce many-body Hilbert space



Work with a subset of many-body wave functions (→ choose simple)



Energy-density functional (EDF) approach

$$H^{eff} |\Psi^{eff}\rangle = E |\Psi^{eff}\rangle$$

$$H^{eff} \neq H^{ab\ initio}$$

parameters phenomenologically adjusted to reproduce nuclear properties

→ loss of predictive power compared to ab initio

→ but simpler calculations and wider reach across the nuclear chart