


Electroweak interaction

The Glashow-Salam-Weinberg model

Particle Physics
Fabrice Couderc / Eli BenHaim

- I. Introduction to $SU(2)_L$
 1. Charged weak current reminder
 2. Building a gauge theory of weak interaction
 3. Gauge bosons masses
 4. Discovery of neutral weak currents
- II. Symmetry breaking and the BEH mechanism
 1. Scalar field and the abelian symmetry case
- III. Electroweak symmetry and GSW model
 1. Electroweak symmetry $SU(2)_L \times U(1)$
 2. Breaking of EWK symmetry and gauge bosons masses
 3. Coupling to fermions, hypercharge and weak isospin
 4. W, Z boson couplings summary
- IV. Experimental tests and exercises
 1. Z boson properties
 2. Lepton coupling universality
 3. Number of neutrino families
 4. Left-Right asymmetry at SLC and $\sin^2\theta_w$

Weak neutral current discovery




Gargamelle bubble chamber 1973

$$\nu_{\mu} e^{-} \rightarrow e^{-} \nu_{\mu}$$

Weak neutral current discovery


Gargamelle bubble chamber 1973



$$\nu_{\mu} p \rightarrow X \nu_{\mu}$$



Z boson discovery

- CERN: UA1 and UA2 located on the SPS (Proton synchrotron) ppbar collider with $\sqrt{s} = 540\text{GeV}$ - 1983



High-energy lepton pair:

$$m_{\ell\ell}^2 = (p_{\ell^+} + p_{\ell^-})^2 = M_Z^2$$





$M_Z \approx 91\text{ GeV}$

- Break symmetry with an SU(2) doublet

$$V(\phi) = m^2 |\phi|^2$$


$$V(\phi) = -\mu^2 |\phi|^2 + \lambda |\phi|^4$$



Developing ϕ around its minimum value $|\phi_{\min}| = v^2/2$

$$v = \sqrt{\frac{\mu^2}{\lambda}}$$


= 246 GeV



Hypercharge and weak isospin

	Q	T ₃ (left)	Y
U	+2/3	+1/2	1/6
D	-1/3	-1/2	1/6
nu	0	+1/2	-1/2
lep	-1	-1/2	-1/2

Can be summarised with
 $Q = T_3 + Y$
 $T_3 (U_L, D_L, \text{right}) = +1/2, -1/2, 0$




Green curve includes radiative corrections

$m_Z = 91.2 \text{ GeV}$
 $\Gamma_Z = 1.5 \text{ GeV}$

- Do these couplings exist $A_\mu \bar{\psi}_L \gamma_\mu \psi_R$, $A_\mu \bar{\psi}_R \gamma_\mu \psi_L$?
 ✓ and for Z_μ ? W_μ ?
- Assuming leptons and quarks are massless,
 - ❖ Compute $B(W \rightarrow \ell \nu)$ at tree level neglecting phase space
 - ❖ Compute $\Gamma(Z \rightarrow \bar{\nu}\nu) / \Gamma(Z \rightarrow \bar{e}e)$
 - ❖ From $B(Z \rightarrow \ell\ell) = 3.3\%$, conclude that $B(Z \rightarrow \nu\nu) \approx 20\%$
 - ❖ Use it to predict the number of neutrino family from Γ_{inv} measurement



Note: $\sin^2\theta_w \approx 0.23$

Number of neutrino family



$$N_\nu = 2.984 \pm 0.008$$

Asymmetry A_{LR} and $\sin^2\theta_w$



$\sin^2\theta_{\text{eff}} = 0.23097 \pm 0.00027$

NB: $1 - (m_w/m_z)^2 = 0.22$