

# Quantum ChromoDynamics

## QCD Lagrangian and SU(3) structure

Particle Physics  
Fabrice Couderc / Eli BenHaim

- QED Lagrangian, reminder
- Building the QCD Lagrangian
  - ✓ Recipe from QED and implications
  - ✓ Feynman rules
- QCD group structure
  - ✓ Gluon emission and gluon splitting
  - ✓ Infra red divergence, soft gluon emission probability
  - ✓ Hadron multiplicity measurements
  - ✓ 3jets / 4 jets measurements
- Color Factor
  - ✓  $qq$  and  $q\bar{q}$  interaction
  - ✓ QCD potential and meson stability
- Strong coupling constant measurements at the LHC
  - ✓ Total cross section at the LHC
  - ✓ Inclusive jet production
  - ✓ 3jets / 2jets ratio

## Refs:

- Daniel Décamp QCD
- Griffiths, chapter 8 (color factors)

Free field propagation (propagators)

Interaction quarks gluon (a la QED)

Vertex 3 gluons ( due to the Abelian group )




Vertex 4 gluons ( due to the Abelian group )

$$\mathcal{L}_{\text{QCD}} = -\frac{1}{4}(\partial_\mu G_\nu^a - \partial_\nu G_\mu^a)(\partial_\mu G_\nu^a - \partial_\nu G_\mu^a) + \bar{\psi}(i\partial - m)\psi$$

$$+ g_s \bar{\psi}_i \gamma^\mu (t^a)_{ij} \psi_j G_\mu^a$$

$$- \frac{g_s}{2} f^{abc} (\partial_\mu G_\nu^a - \partial_\nu G_\mu^a) G^{b\mu} G^{c\nu}$$

$$- \frac{g_s^2}{4} f^{abc} f^{ade} G_\mu^b G_\nu^c G^{d\mu} G^{e\nu}$$



Fermions lines carry a color charge!


$$c_i u_f^{(s)}(p^\mu) \text{ incoming}$$


$$c_i^\dagger \bar{u}_f^{(s)}(p^\mu) \text{ outgoing}$$


$$c_i^\dagger \bar{v}_f^{(s)}(p^\mu) \text{ incoming}$$

$$c_i v_f^{(s)}(p^\mu) \text{ outgoing}$$

Interaction (or completeness)

Fermion vertex:  =  $ig\gamma^\mu t^a$

3-boson vertex:  =  $gf^{abc}[g^{\mu\nu}(k-p)^\rho + g^{\nu\rho}(p-q)^\mu + g^{\rho\mu}(q-k)^\nu]$

4-boson vertex:  =  $-ig^2[f^{abe}f^{cde}(g^{\mu\rho}g^{\nu\sigma} - g^{\mu\sigma}g^{\nu\rho}) + f^{ace}f^{bde}(g^{\mu\nu}g^{\rho\sigma} - g^{\mu\sigma}g^{\nu\rho}) + f^{ade}f^{bce}(g^{\mu\nu}g^{\rho\sigma} - g^{\mu\rho}g^{\nu\sigma})]$

One trivial basis for color


$$c_r = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad c_g = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad c_b = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

$$\begin{aligned}
 t^1 &= \frac{1}{2} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & t^2 &= \frac{1}{2} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & t^3 &= \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\
 t^4 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix}, & t^5 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix}, \\
 t^6 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, & t^7 &= \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix}, & t^8 &= \frac{1}{2\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}.
 \end{aligned}$$

# The group structure of QCD

## From the PDG

Useful color-algebra relations include:  $t_{ab}^A t_{bc}^A = C_F \delta_{ac}$ , where  $C_F \equiv (N_c^2 - 1)/(2N_c) = 4/3$  is the color-factor (“Casimir”) associated with gluon emission from a quark;  $f_{ACD} f_{BCD} = C_A \delta_{AB}$  where  $C_A \equiv N_c = 3$  is the color-factor associated with gluon emission from a gluon;  $t_{ab}^A t_{ab}^B = T_R \delta_{AB}$ , where  $T_R = 1/2$  is the color-factor for a gluon to split to a  $q\bar{q}$  pair.



$$\sum_{a=1}^{N_A} (T^a T^{\dagger a})_{ij} = \delta_{ij} C_F$$

$$\sum_{a,b=1}^{N_A} f^{abc} f^{*abd} = \delta^{cd} C_A$$


$$\sum_{i,j=1}^{N_F} T_{ij}^a T_{ji}^{\dagger b} = \delta^{ab} T_F$$

# Measuring the group structure

$N_{ch}$  = charge hadron multiplicity in a jet



Ratio of the “slopes”

$$R = 2.22 \pm 0.11$$




# Measuring the group structure





- LEP events with 3 and 4 jets





# Quark - antiquark interaction






## Example of qq and qqbar scattering


$q\bar{q} \rightarrow q\bar{q}$	$qq \rightarrow qq$	$f(ijkl)$
$xx \rightarrow xx$	$xx \rightarrow xx$	$f(xxxx) = +1/3$
$xx \rightarrow yy$	$xy \rightarrow yx$	$f(yxxy) = +1/2$
$xy \rightarrow xy$	$xy \rightarrow xy$	$f(xxyy) = -1/6$
$xy \rightarrow yx$	$xx \rightarrow yy$	$f(yxyx) = 0$

# QCD running measurement

Measuring the running of  $\alpha_s$  is still a very active field at LHC  
 Important to predict any cross section!!!



# Total cross section at the LHC



# R32 : 3jets to 2jets production

Naively  $\sigma[2 \text{ jets}] \propto \alpha_s^2$  ;  $\sigma[3 \text{ jets}] \propto \alpha_s^3 \Rightarrow \sigma[3 \text{ jets}] / \sigma[2 \text{ jets}] \propto \alpha_s$

several systematics cancel in the ratio

(e.g. pdf uncertainty greatly reduced)

$$\mu_R = (p_{T1} + p_{T2}) / 2$$

World average:

$$\alpha_s(M_Z) = 0.1184 \pm 0.0007$$

CMS 3/2 jets:

$$\alpha_s(M_Z) = 0.1148 \pm 0.006$$

