

Exercise sheet № 6 - QCD: color factors and parton density functions

1 Color factors

We consider $q\bar{q}'$ scattering at low energy. In this case the classical potential between the pair quark anti-quark is the classical limit of the matrix element:

$$V_{strong} = \langle q\bar{q}' | \mathcal{M}_{int} | q\bar{q}' \rangle$$

where \mathcal{M} is the matrix element of the scattering diagram.

1. Draw the Feynman diagram of the scattering $q\bar{q}'$ considering only the strong interaction.
2. Find out that the classical potential $V_{q\bar{q}'} = F_{col}\alpha_w V_{EM}$ where V_{EM} is the classical electromagnetic potential (without the hyperfine structure constant α_{EM}). Give the expression of the color factor F_{col} as a function f_{ijkl} the color factors defined in the course. In this case the two partons have an opposite color charge (color vs anti-color) therefore V_{EM} is an attractive potential unless F_{col} is negative.
3. We consider $q\bar{q}'$ state in the "colorless" configuration of the octet, *i.e.*

$$|8\rangle = \frac{1}{\sqrt{6}} |r\bar{r} + g\bar{g} - 2b\bar{b}\rangle$$

Find the color factor F_{col} for the scattering $|8\rangle \rightarrow |8\rangle$. What is its sign? What do you conclude on the stability of this "colorless" state.

4. What color factor do you expect for the scattering $|8\rangle \rightarrow |0\rangle$, where

$$|0\rangle = \frac{1}{\sqrt{3}} |r\bar{r} + g\bar{g} + b\bar{b}\rangle$$

is the singlet meson state? Compute this color factor and conclude.

5. Eventually find the color factor of the scattering $|0\rangle \rightarrow |0\rangle$? What is its sign and what does it mean?

2 Nucleon momentum

We note f_q^p (resp. f_q^n) the probability density function of parton q in the proton (resp. in the neutron). In this exercise we will only consider high x and low Q^2 therefore limit ourselves to valence quarks and gluons.

1. Write the total momentum of the proton as a function of the integrals of f_u^p , f_d^p and f_g^p . We note $\epsilon_q^p = \int x f_q^p(x) dx$, find out the conservation rule:

$$\epsilon_u^p + \epsilon_d^p + \epsilon_g^p = 1$$

2. Using the fact that strong interaction conserves isospin, write the same expression for the neutron, and simplify your expression using the notation for all partons q

$$\epsilon_q^p \equiv \epsilon_q$$

3. We remind you that the form factor of the proton involved in photon-nucleon scattering is $F_2^p(x) = \sum_q Q_q^2 f_q^p(x)$ where Q_q is the charge of parton q in unit of the electron charge. Write the expression of $\int x F_2^p(x) dx$ as a function of ϵ_q 's
4. Similarly for the neutron, we define $F_2^n(x) = \sum_q Q_q^2 f_q^n(x)$. Write this integral as a function of ϵ_q 's and use again the isospin to get the expression as a function of the ϵ_q 's
5. Experimentally, we measured:

$$\begin{aligned} \int_0^1 x F_2^p(x) dx &= 0.18 , \\ \int_0^1 x F_2^n(x) dx &= 0.12 . \end{aligned} \tag{1}$$

From these two measurements conclude on the value of ϵ_g the fraction of the nucleon momentum carried out by gluons. This was one of the first evidence for the presence in the nucleon of something else than valence quarks.